Contents

Conference Photograph vi

Preface vii

Conference Information ix

Chapter 1. Fractional quantum Hall effect

1.1. Composite particles and skyrmions

Skyrmion excitations in the limit of vanishing Landé g-factor
D.K. Maude, S. Marty, L.B. Rigal, M. Potemski, J.C. Portal, Zb. Wasilewski, M. Henini, L. Eaves, G. Hill and M.A. Pate 1

Absorption spectroscopy of charged spin texture excitations at $\nu = 1$
B.B. Goldberg, M.J. Manfra, L. Pfeiffer and K. West 7

Composite fermions in magnetic focusing and commensurability experiments
J.H. Smet, D. Weiss, K. von Klitzing, R. Fleischmann, R. Ketzmerick, T. Geisel, W. Wegscheider, P.T. Coleridge, Z.W. Wasilewski and G. Weimann 15

Measurements of composite Skyrmions at filling factor $\frac{1}{3}$
D.R. Leadley, R.J. Nicholas, D.K. Maude, A.N. Utjuzh, J.C. Portal, J.J. Harris and C.T. Foxon 23

Manifestation of charge-flux duality in a 2D electron gas near filling factor $\nu = \frac{1}{2}$
G.M. Gusev, Z.D. Kvon, E.B. Olshanetsky, D.K. Maude, X. Kleber and J.C. Portal 27

Fermi-liquid shell model for N electrons on a spherical surface: A test of the composite fermion hierarchy picture
P. Sitko, K.-S. Yi and J.J. Quinn 32

Composite fermion signature in the acoustoelectric effect in 2DES and 2DHS
I. Kennedy, V.W. Rampton, C.J. Mellor, B.H. Bracher, M. Henini, Z.R. Wasilewski and P.T. Coleridge 36

1.2. Fractional quantum Hall liquid and Wigner solid

Light scattering by collective excitations in the fractional quantum Hall regime
A. Pinczuk, B.S. Dennis, L.N. Pfeiffer and K.W. West 40

Role of spin excitations in the fractional quantum Hall effect at $\nu = \frac{1}{3}$
J.C. Harris, H.D.M. Davies, J.F. Ryan and A.J. Turberfield 44

Ballistic phonon absorption in the fractional and non-quantised Hall effects
U. Zeitler, A.M. Devitt, J.E. Digby, C.J. Mellor, A.J. Kent, K. Benedict and T. Cheng 49

Microwave absorption in the magnetically-induced Wigner solid phase of a two-dimensional hole system
P.F. Hennigan, A. Beya, C.J. Mellor, R. Gaál, F.I.B. Williams and M. Henini 53

Microwave conductivity of 2D hole systems at low filling
C.-C. Li, L.W. Engel, D. Shahar, D.C. Tsui and M. Shayegan 57

Magnetism in a Wigner solid and Aharonov–Bohm effect
T. Okamoto and S. Kawaji 61

Effect of valence band structure on the fractional quantum Hall effect of two-dimensional hole systems
K. Muraki and Y. Hirayama 65

Chapter 2. Integer quantum Hall effect

2D transport at the surface of a 3D quantum Hall system
D.P. Druist, P.J. Turley, E.G. Gwinn, K. Maranowski and A.C. Gossard 70

The quantum Hall effect and chiral Fermi liquids in a quasi-two-dimensional organic conductor at very high magnetic fields
H. Yaguchi, N. Harrison, M.M. Honold, C. Mielke, J. Singleton, P.J. Gee, D. Rickel, I. Deckers, P.H.P. Reinders, F. Herlach, M. Kurmoo and P. Day 75

Scanned potential microscopy of a two-dimensional electron gas
K.L. McCormick, M.T. Woodside, M. Huang, P.L. McEuen, C.I. Duruoz and J.S. Harris Jr 79

Quantum-to-classical crossover of quantum Hall effect
T. Ando 84

QHE breakdown induced by a single antidot: Coulomb staircase in the breakdown current
Z.H. Liu, G. Nachtwei, J. Weis, K. von Klitzing and K. Eberl 89

Breakdown of the integer quantum Hall effect studied by Corbino discs
M. Yokoi, T. Okamoto, S. Kawaji, T. Goto and T. Fukase 93

Bulk current distribution in Hall bar geometry samples in the integer quantum Hall effect
E. Yahel, A. Palevski and H. Shtrikman 97

Temperature dependence of the current distribution in the quantum hall regime in a distributed magnetic field
J. Wakabayashi, A. Tamagawa, T. Nagashima, T. Mochiku and K. Hirata 102

Hall electric field-dependent broadening of extended state bands in Landau levels and breakdown of the quantum Hall effect
T. Shimada, T. Okamoto and S. Kawaji 107

Heat capacity and nuclear-spin dynamics in GaAs/AlGaAs heterostructures around $\nu = 1$
V. Bayot, E. Grivei, S. Melinte and M. Shayegan 111

Magnetization and incompressibility of a high mobility 2D electron gas in the IQHE
S.A.J. Wiegers, M. Specht, L.P. Lévy, M.Y. Simmons, D.A. Ritchie, A. Cavanna, B. Etienne, G. Martinez and P. Wyder 115

Millimeterwave conductivity experiments regarding dynamical scaling of the integer quantum Hall effect
W. Belitsch, R. Meisels, F. Kuchar, G. Hein and K. Pierz 119

Activated conductivities and nonuniversal behaviour in large high mobility Hall bars
M. Furlan 123

Phase coherence of edge states over macroscopic length scales
T. Machida, H. Hirai, S. Komiyama and Y. Shiraki 128

Chapter 3. Quantum wires

Transport spectroscopy of single-walled carbon nanotubes
D.H. Cobden, M. Bockrath, N.G. Chopra, A. Zettl, P.L. McEuen, A. Rinzler, A. Thess and R.E. Smalley 132

Scattering by topological disorder in connected carbon nanotubes in magnetic fields
T. Nakanishi and T. Ando 136

Quantized current in one-dimensional channel induced by surface acoustic waves
V.I. Talyanskii, J.M. Shilton, J. Cunningham, M. Pepper, C.B. Ford, C.G. Smith, E.H. Linfield, D.A. Ritchie and G.A.C. Jones 140

Giant oscillations of acousto-conductance in a quantum channel
H. Totland, Ø.L. Bø and Y.M. Galperin 147

Signs of quantum statistical effects in electron collision
R.C. Liu, Y. Yamamoto and S. Tarucha 152

Closely separated one-dimensional wires: coupled ballistic conduction, wave function hybridization and compressibility investigations
I.M. Castleton, A.G. Davies, A.R. Hamilton, J.E.F. Frost, M.Y. Simmons, D.A. Ritchie and M. Pepper 157

Size-effects and tuning of phase coherence in a double quantum well mesoscopic wire
M.J. Gompertz, T. Ihn, P.C. Main, A. Nogaret, L. Eaves, M. Henini and S.P. Beaumont 162

Multiple subband crossing in a one-dimensional hole gas with enhanced g-factors
A.J. Daneshvar, C.J.B. Ford, A.R. Hamilton, M.Y. Simmons, M. Pepper and D.A. Ritchie 166

Polarization anisotropy and Zeeman splitting in strained quantum wires: Dimensional crossover effect on strain
M. Notomi, J. Hammersberg, J. Zeman, H. Weman, M. Potemski, H. Sugiura and T. Tamamura 171

Experimental determination of spectral densities in quasi-one-dimensional electron systems
B. Kardynal, C.H.W. Barnes, E.H. Linfield, D.A. Ritchie, J.T. Nicholls, K.M. Brown, C.J.B. Ford and M. Pepper 175

Temperature dependence of the "0.7" $2e^2/h$ quasi-plateau in strongly confined quantum point contacts
A. Kristensen, P.E. Lindelof, J. Bo Jensen, M. Zaffalon, J. Hollingbery, S.W. Pedersen, N. Nygård, H. Bruus, S.M. Reimann, C.B. Sørensen, M. Michel and A. Forchel 180

Optical detection of critical exponents in the Tomonaga–Luttinger liquid
T. Ogawa 185

Chapter 4. Quantum dots

Coulomb oscillations in few-electron quantum dot
L.P. Kouwenhoven, T.H. Oosterkamp, S. Tarucha, D.G. Austing and T. Honda 191

Micro-mechanical charge transfer mechanism in soft Coulomb blockade nanostructures
L.Y. Gorelik, A. Isacsson, M. Jonson, B. Kasemo, R.I. Shekhter and M.V. Voinova 197

Statistics of peak spacings and widths in the quantum Coulomb blockade regime
C.M. Marcus, S.R. Patel, C.I. Duruöz, J.S. Harris Jr., K. Campman and A.C. Gossard 201

Quantum dot molecules
D.G. Austing, T. Honda, K. Muraki, Y. Tokura and S. Tarucha 206

Numerical study of spin effects in single and double quantum dots
H. Tamura 210

Symmetry of 'molecular' configurations of interacting electrons in a quantum dot in strong magnetic fields
H. Imamura, P.A. Maksym and H. Aoki 214

Probing of level correlation effects in quantum dots by surface acoustic-waves
A. Knäbchen, O. Entin-Wohlman and Y.B. Levinson 220

Effect of interactions on the integrability and level statistics of two particles in an infinite quantum well
L. Meza-Montes, and S.E. Ulloa 224

Fluctuations in quantum dot charging energy
M. Stopa 228

Quantum states of interacting electrons in a 2D elliptical quantum dot
P.A. Maksym 233

Electronic structures in a quantum dot with asymmetric confining potential
T. Ezaki, Y. Sugimoto, N. Mori and C. Hamaguchi 238

Resonant tunneling through coupled InAs quantum dots
K. Yoh, H. Kazama and T. Nakano 243

Magneto-photoluminescence in high magnetic fields from InGaAs/GaAs quantum dots formed in tetrahedral-shaped recesses
K. Uchida, N. Miura, Y. Sakuma, Y. Awano, T. Futatsugi and N. Yokoyama 247

Coulomb blockade in arrays of quantum dots
M. Hirasawa, S. Katsumoto, A. Endo and Y. Iye 252

Fine structure in the spectrum of the few-electron ground states of self-assembled quantum dots
B.T. Miller, W. Hansen, S. Manus, R.J. Luyken, A. Lorke, J.P. Kotthaus, G. Medeiros-Ribeiro and P.M. Petroff 257

Photoluminescence spectroscopy of self-assembled (InGa)As quantum dots in high magnetic fields
R.K. Hayden, K. Uchida, N. Miura, A. Polimeni, S.T. Stoddart, M. Henini, L. Eaves and P.C. Main 262

Electron escape from self-assembled InAs/GaAs quantum dot stacks
P.N. Brounkov, A.A. Suvorova, M.V. Maximov, A.F. Tsatsul'nikov, A.E. Zhukov, A.Yu. Egorov, A.R. Kovsh, S.G. Konnikov, T. Ihn, S.T. Stoddart, L. Eaves and P.C. Main 267

Capacitance of InAs quantum dot arrays in a mean field approach
O. Heller, Ph. Lelong and G. Bastard 271

Coulomb blockade and enhancement of magnetoresistance change in ultrasmall ferromagnetic tunnel junctions
S. Iwabuchi, T. Tanamoto and R. Kitawaki 276

Thermopower measurements of semiconductor quantum dots
A.S. Dzurak, C.G. Smith, C.H.W. Barnes, M. Pepper, L. Martín-Moreno, C.T. Liang, D.A. Ritchie and G.A.C. Jones 281

Plasmons in spatially non-uniform magnetic fields
S. Cina, D.D. Arnone, J.H. Burroughes, S.N. Holmes, T. Burke, H.P. Hughes, D.A. Ritchie and M. Pepper 286

Edge state formation in magnetic quantum structures
G. Ihm, N. Kim, H.-S. Sim, K.-H. Ahn, K.J. Chang and S.J. Lee 291

Chapter 5. Ballistic transport

Dephasing due to *which path* detector
E. Buks, R. Schuster, M. Heiblum, D. Mahalu and V. Umansky 295

Insulator–quantum Hall liquid transition in antidot lattices
F. Nihey, K. Nakamura, M.A. Kastner, T. Takamasu and G. Kido 302

Localization oscillation in antidot lattices
S. Uryu and T. Ando 308

Far-infrared and transport properties of antidot arrays with broken symmetry
A. Lorke, S. Wimmer, B. Jager, J.P. Kotthaus, W. Wegscheider and M. Bichler 312

Manifestation of commensurate orbits in the magnetoluminescence spectrum of electrons in antidot arrays
I.V. Kukushkin, D. Weiss, G. Lütjering, R. Bergmann, H. Schweizer, K.V. Klitzing, K. Eberl, P. Rotter, M. Suhrke and U. Rössler 317

Negative magnetoresistance in antidot lattices with saddle points
R. Henning, P. Rotter, M. Suhrke and U. Rössler 321

Giant magnetoresistance and possible miniband effects in periodic magnetic fields
N. Overend, A. Nogaret, B.L. Gallagher, P.C. Main, R. Wirtz, R. Newbury, M.A. Howson and S.P. Beaumont 326

Huge magnetoresistance oscillations in periodic magnetic fields
P.D. Ye, D. Weiss, R.R. Gerhardts, K. von Klitzing and S. Tarucha 330

Self-similar conductance fluctuations in a Sinai billiard with a mixed chaotic phase space
T.M. Fromhold, C.R. Tench, R.P. Taylor, A.P. Micolich and R. Newbury 334

Frequency-dependent magnetotransport in a two-dimensional magnetic modulation system
E. Badran and S.E. Ulloa 339

Geometry-induced fractal behaviour: fractional Brownian motion in a ballistic mesoscopic billiard
A.P. Micolich, R.P. Taylor, R. Newbury, J.P. Bird, R. Wirtz, T.M. Fromhold, Y. Aoyagi and T. Sugano 343

Dephasing in open quantum dots
A.G. Huibers, M. Switkes, C.M. Marcus, K. Campman and A.C. Gossard 348

Wave function scarring and magnetotransport in quantum dots
Y. Ochiai, Y. Okubo, N. Sasaki, J.P. Bird, K. Ishibashi, Y. Aoyagi, T. Sugano, A.P. Micolich, R.P. Taylor, R. Newbury, D. Vasileska, R. Akis and D.K. Ferry 353

Wave function scarring effects in Sinai billiard cavities
R. Akis and D.K. Ferry 358

Resonant tunnelling spectroscopy of a two-dimensional 'tear-shaped' billiard
T.M. Fromhold, A. Nogaret, P. Hawrylak, C.R. Tench, F.W. Sheard, L. Eaves and P.C. Main 364

Stability of regular orbits in ballistic quantum dots
R. Akis, D. Vasileska, D.K. Ferry, J.P. Bird, Y. Okubo, Y. Ochiai, J.P.K. Ishibashi, Y. Aoyagi and T. Sugano 368

Dynamic barriers in uniform 2DEG as splitter and source of electron beams
O.A. Tkachenko, D.G. Baksheyev, M.B. Wojtsekhowski and V.A. Tkachenko 373

Effect of curvature of a 2D electron sheet on the ballistic conductance and spin–orbit interaction
A.V. Chaplik, L.I. Magarill and D.A. Romanov 377

Chapter 6. Edge states and nonequilibrium properties

An apparent $g = \frac{1}{2}$ chiral Luttinger liquid at the edge of the $\nu = \frac{1}{2}$ compressible composite Fermion liquid
A.M. Chang, L.N. Pfeiffer and K.W. West 383

Coherent transport through a quantum dot in a strong magnetic field
M.R. Geller 388

Possible explanation of the high temperature extrapolated value of diagonal resistivity at $\nu = 1$ in terms of skyrmions
T. Takamasu, G. Kido, M. Ohno, N. Miura, A. Endo, M. Kato, S. Katsumoto and Y. Iye 391

Direct observation of a fractional charge
R. de-Picciotto, M. Reznikov, M. Heiblum, V. Umansky, G. Bunin and D. Mahalu 395

Observation of $e/3$ fractionally charged quasiparticles
D.C. Glattli, L. Saminadayar, Y. Jin and B. Etienne 401

Non-equilibrium transport along an edge of variable slope in the fractional quantum Hall regime
J.D.F. Franklin, C.J.B. Ford, M.Y. Simmons, D.A. Ritchie and M. Pepper 405

Tunneling through point contacts in the quantum Hall effect
P.J. Turley, D.P. Druist, E.G. Gwinn, K. Maranowski, K. Campmann and A.C. Gossard 410

Self-consistent edge states of quantum wires in high magnetic fields
T. Suzuki and T. Ando 415

Quantum transport in fractional quantum Hall edges
K.-i. Imura and N. Nagaosa 420

Conduction through point contacts in fractional quantum Hall liquid
M. Ando, A. Endo, S. Katsumoto and Y. Iye 426

Resonant tunneling between quantum Hall edges through a quantum dot
A. Furusaki 430

High frequency edge excitations in the quantum Hall regime
N.Q. Balaban, U. Meirav, H. Shtrikman and V. Umansky 435

Edge channels in integer and fractional quantum-Hall regime investigated by magnetocapacitance
K. Oto, S. Uno, S. Takaoka, K. Murase, F. Nihey and K. Nakamura 440

Chapter 7. SN junctions

Mesoscopic superconductors as 'artificial atoms' made from Cooper pairs
A.K. Geim, I.V. Grigorieva, S.V. Dubonos, J.G.S. Lok, J.C. Maan, A.E. Filippov, F.M. Peeters and P.S. Deo 445

Fluxoid states in mesoscopic superconductors
H. Sato, S. Katsumoto and Y. Iye 453

Energy spectroscopy of Andreev levels between two superconductors
A.F. Morpurgo, B.J. van Wees, T.M. Klapwijk and G. Borghs 458

Semiconductor-coupled superconducting junctions using NbN electrodes with high H_{c2} and T_c
H. Takayanagi and T. Akazaki 462

Reentrant resistance and giant Andreev backscattering in a two-dimensional electron gas coupled to superconductors
S.G. den Hartog, B.J. van Wees, Yu.V. Nazarov, T.M. Klapwijk and G. Borghs 467

Magnetoresistance oscillations of quasi-ballistic two-dimensional electron gas coupled with a superconducting ring
E. Toyoda and H. Takayanagi 472

Phase boundaries of superconducting mesoscopic square loops
V.M. Fomin, V.R. Misko, J.T. Devreese and V.V. Moshchalkov 476

A new singularity in the coherent coupling in Al/GaAs/Al SNS junctions at the bias voltage corresponding to the superconducting energy gap
R. Taboryski, J. Kutchinsky, O. Kuhn, P. Lindelof, C.B. Sørensen, A. Kristensen, C.S. Jacobsen, J.B. Hansen and J.L. Skov 480

Sample-specific conductance fluctuations modulated by the superconducting phase
S.G. den Hartog, C.M.A. Kapteyn, B.J. van Wees, T.M. Klapwijk and G. Borghs 485

Collective charge density wave motion in inhomogeneous mesoscopic systems
M.I. Visscher, B. Rejaei and G.E.W. Bauer 490

Chapter 8. Mesoscopic effects

Single-electron transistor as an electrometer on a two-dimensional electron system
Y.Y. Wei, J. Weis, K.V. Klitzing and K. Eberl 496

Probing spin dynamics by conductance fluctuations and noise in mesoscopic spin-glass
J. Jaroszyński, T. Dietl, J. Wróbel, G. Karczewski, T. Wojtowicz, G. Grabecki, M. Sawicki, E. Papis, E. Kamińska and A. Piotrowska 500

Conductance fluctuations near the two-dimensional metal–insulator transition
K.P. Li, D. Popović and S. Washburn 504

Experiments on Aharonov–Bohm effect under the influence of uniform spin–orbit interaction: possible observation of Berry's phase in electronic transport
A.F. Morpurgo, J.P. Heida, B.J. van Wees, T.M. Klapwijk and G. Borghs 509

Quantum transport in ferromagnetic dot structure embedded in semiconductor quantum wires
T. Kikutani, N. Aoki, C.U. Hong, H. Hori and S. Yamada 513

Interacting one-dimensional electrons driven by two-dimensional reservoir electrons
A. Shimizu and T. Miyadera 518

Quasi-ballistic conductance fluctuations and ergodic hypothesis
Y. Asano and G.E.W. Bauer 523

Chapter 9. Optical properties

Terahertz dynamics in confined magnetoexcitons
J. Kono, M.Y. Su, J. Černe, M.S. Sherwin, S.J. Allen Jr, T. Inoshita, T. Noda and H. Sakaki 527

Mixing of terahertz and near-infrared radiation in quantum wells in strong magnetic fields
T. Inoshita and H. Sakaki 534

Magneto-optical study of excitonic states in 2DEGs near filling factor $\nu = 1$
J.L. Osborne, A.J. Shields, M.Y. Simmons, N.R. Cooper, D.A. Ritchie and M. Pepper 538

Skyrmion–hole excitations at $\nu = 1$ studied by photoluminescence spectroscopy
H.D.M. Davies, J.C. Harris, R.L. Brockbank, J.F. Ryan, A.J. Turberfield, M.Y. Simmons and D.A. Ritchie 544

Photoluminescence in quantum Hall regime: Exact diagonalization study
K. Asano and T. Ando 549

Photoluminescence discontinuities close to the filling factor $\nu = 1$
R.J. Nicholas, D. Kinder, A.N. Priest, C.C. Chang, H.H. Cheng, J.J. Harris and C.T. Foxon 553

Spin-dependent photoluminescence at $\nu = 1$ filling factor
F. Plentz, D. Heiman, A. Pinczuk, L.N. Pfeiffer and K.W. West 558

Photoluminescence of GaAs/AlGaAs heterojunctions in magnetic fields up to 50 T
A.N. Priest, R.J. Nicholas, H.H. Cheng, M. Van der Burgt, J.J. Harris and C.T. Foxon 562

Emission from a two-dimensional electron gas in quantizing magnetic fields
M. Potemski, L. Gravier, A. Trüby, D.K. Maude, P. Hawrylak, Zb. Wasilewski and B. Etienne 566

Negative vertical photovoltaic response of two-dimensional electrons in the quantum Hall regime
S. Tsujino, H. Nakajima, T. Inoshita, T. Noda and H. Sakaki 571

Shake-up processes of a two-dimensional electron gas in GaAs/AlGaAs quantum wells at high magnetic fields
G. Finkelstein, H. Shtrikman and I. Bar-Joseph 575

Near surface quantum well excitons in magnetic fields
S.G. Tikhodeev, N.A. Gippius, A.B. Dzyubenko, L.V. Kulik, V.D. Kulakovskii and A. Forchel 580

Charged excitons under applied electric and magnetic fields
A.J. Shields, J.L. Osborne, D.M. Whittaker, M.Y. Simmons, F.M. Bolton, D.A. Ritchie and M. Pepper 584

Difference of self-energy corrections for charged and uncharged states in degenerate two dimensional systems
J.X. Shen, Y. Oka, W. Ossau, F. Fischer, A. Waag and G. Landwehr 589

Exciton and charged density wave formation in spatially separated electron–hole liquids
L. Liu, L. Świerkowski and D. Neilson 594

Internal transitions of excitons and hole cyclotron resonance in undoped GaAs/AlGaAs quantum wells by optically detected resonance spectroscopy
H.A. Nickel, G.S. Herold, M.S. Salib, G. Kioseoglou, A. Petrou, B.D. McCombe and D. Broido 598

Role of the valence-band mass on the many-body optical properties of a two-dimensional electron gas
H.P. van der Meulen, J. Rubio, J.M. Calleja, C. Tejedor, F. Rodriguez, V. Haerle and F. Scholz 603

Interband spectroscopy of p-type GaAs/(Al,Ga)As quantum wells at low hole densities
B.E. Cole, T. Takamasu, K. Takehana, R. Goldhahn, D. Schulze, G. Kido, J.M. Chamberlain, G. Gobsch, M. Henini and G. Hill 607

A quantum Monte Carlo study on excitonic molecules in type-II superlattices
T. Tsuchiya and S. Katayama 612

Manifestation of the magnetic depopulation of one-dimensional subbands in the optical absorption of acoustic magnetoplasmons in side-gated quantum wires
C. Steinebach, T. Kurth, D. Heitmann and V. Gudmundsson 617

Exciton complexes in InGaAs/GaAs quantum dots
M. Bayer, V.D. Kulakovskii, T. Gutbrod and A. Forchel 620

Magnetic trap for excitons
P.C.M. Christianen, F. Piazza, J.G.S. Lok, J.C. Maan and W. van der Vleuten 624

Chapter 10. Electrons on liquid helium

Magnetotransport in the two-dimensional electron fluid and solid and liquid helium
M.J. Lea and M.I. Dykman 628

Wigner solid dynamics on normal and superfluid ^{3}He
Yu.P. Monarkha, K. Shirahama, O.I. Kirichek and K. Kono 636

Phase diagram of the screened two-dimensional electron crystal
G. Mistura, S. Neser and P. Leiderer 640

Polaron transition of surface electrons on liquid ^{4}He films in zero and quantizing magnetic fields
O. Tress, F.C. Penning, G. Mistura, E. Teske and P. Wyder 644

Mobility of a 2D Wigner crystal over liquid ^{4}He and ^{3}He–^{4}He solution
V.V. Dotsenko, V.E. Sivokon, Yu.Z. Kovdrya and V.N. Grigor'ev 648

Observation of magnetoplasma resonances in a single wire of electrons on helium
A.M.C. Valkering and R.W. van der Heijden 652

Conduction of electrons on liquid helium along channels produced by multi-layer microfabrication
R.J.F. van Haren, G. Acres, P. Fozooni, A. Kristensen, M.J. Lea, P.J. Richardson,
A.M.C. Valkering and R.W. van der Heijden 656

Contact phenomena in 2D electron systems
V. Shikin, E. Teske, P. Leiderer and P. Wyder 660

Modes of transverse response in a two-dimensional Coulomb system above the melting temperature
P.L. Elliott, C.I. Pakes, L. Skrbek and W.F. Vinen 664

Damage and annealing in two-dimensional Coulomb crystals
P.L. Elliott, C.I. Pakes, L. Skrbek and W.F. Vinen 668

Mobility of surface electrons over the ^{3}He–^{4}He microstratified solution
S.S. Sokolov, J.-P. Rino and N. Studart 672

Chapter 11. Heterostructure

Spin transport and optically-probed coherence in magnetic semiconductor heterostructures
I.P. Smorchkova, F.S. Flack, N. Samarth, J.M. Kikkawa, S.A. Crooker and D.D. Awschalom 676

Zero-field spin splitting in GaAs/AlGaAs 2D holes
J.P. Lu and M. Shayegan 685

Many-body effects in a quantised 2DES probed by discrete-level tunnelling spectroscopy
A.S.G. Thornton, T. Ihn, P.C. Main, L. Eaves, K.A. Benedict and M. Henini 689

Magnetic properties of a spin-polarized two-dimensional electron system
D. Grundler, I. Meinel, S. Bargstädt-Franke and D. Heitmann 693

Breakdown of the anomalous two-dimensional metallic phase in a parallel magnetic field
V.M. Pudalov, G. Brunthaler, A. Prinz and G. Bauer 697

Metal–insulator transition in a low-mobility two-dimensional electron system
D. Popović, A.B. Fowler and S. Washburn 701

Metal–insulator transition at $B = 0$ in an ultra-low density two-dimensional hole gas
M.Y. Simmons, A.R. Hamilton, T.G. Griffiths, A.K. Savchenko, M. Pepper and D.A. Ritchie 705

Composite electron–hole states in GaSb/InAs quantum wells
A.R. Rundell, G.P. Srivastava, J.C. Inkson, E.A. Johnson and A. MacKinnon 710

Condensation of 2D electron-hole pairs with spin degree of freedom
J. Fernández-Rossier and C. Tejedor 714

Phonon emission by hot 2D electrons and holes in gallium arsenide heterojunction: The influence of screening and acoustic anisotropy
D. Lehmann, Cz. Jasiukiewicz and A.J. Kent 718

Transient negative photocondutance in a charge transfer double quantum well under optical intersubband excitation
M. Rüfenacht, S. Tsujino and H. Sakaki 723

Coupled spin and charge collective excitations in a spin-polarized electron gas
D.C. Marinescu, J.J. Quinn and K.S. Yi 727

Subband structure and anomalous valley splitting in ultra-thin silicon-on-insulator MOSFET's
T. Ouisse, D.K. Maude, S. Horiguchi, Y. Ono, Y. Takahashi, K. Murase and S. Cristoloveanu 731

Electron confinement in (ordered)$GaInP_2$/GaAs/(ordered)$GaInP_2$ single quantum well
J. Zeman, G. Martinez, S.H. Kwok, P.Y. Yu and K. Uchida 735

Magnetophonon and magneto-intersubband-scattering effects in InAs/AlGaSb heterostructures
S. Osako, T. Hamano, N. Mori, C. Hamaguchi, S. Sasa and M. Inoue 740

Magnetothermopower of double p-type quantum wells
R.J. Hyndman, S.T. Stoddart, B. Tieke, S.G.S. Lok, B.L. Gallagher, A.K. Geim, J.C. Maan and M. Henini 745

Interaction ruled temperature dependence of the electron cyclotron mass in GaAs heterojunctions
C.M. Hu, E. Batke, K. Köhler and P. Ganser 749

Strain-induced potential modulation versus magnetic field modulation on two-dimensional electron gas at the GaAs/AlGaAs heterointerface
M. Kato, A. Endo, S. Katsumoto and Y. Iye 753

Suppression of the localization in the two-dimensional conductivity of Si-δ-doped GaAs by Mn co-doping
K.-J. Friedland, I.S. Shlimak and P. Schützendübe 758

Cyclotron resonance of interacting quantum Hall droplets
M. Widmann, U. Merkt, M. Cortés, W. Häusler and K. Eberl 762

Dynamical studies of the radiative recombination process in modulation doped GaAs/AlGaAs heterostructure
T. Lundström, P.O. Holtz, J.P. Bergman, A. Buyanov, B. Monemar, K. Campman, J.L. Merz and A.C. Gossard 767

Spin-splitting collapse in a 2D electron gas in AlGaAs/GaAs heterostructure
G.M. Gusev, Z.D. Kvon, E.B. Olshanetsky, D.K. Maude and J.C. Portal 771

Low-temperature conduction and giant negative magnetoresistance in III–V-based diluted magnetic semiconductor: (Ga, Mn)As/GaAs
A. Oiwa, S. Katsumoto, A. Endo, M. Hirasawa, Y. Iye, F. Matsukura, A. Shen, Y. Sugawara and H. Ohno 775

Distortion of Fermi surface induced by parallel magnetic field in a two-dimensional electron gas
K. Ohtsuka, S. Takaoka, K. Oto, K. Murase and K. Gamo 780

Freely suspended two-dimensional electron gases
R.H. Blick, M.L. Roukes, W. Wegscheider and M. Bichler 784

Theory of AC conductivity of 2D Wigner solids
M. Saitoh 788

Energy-level statistics and localization of 2d electrons in random magnetic fields
M. Batsch, L. Schweitzer and B. Kramer 792

Disordered critical wave functions of two-dimensional Dirac fermions on a lattice
Y. Hatsugai, Y. Morita, X.-G. Wen and M. Kohmoto 796

Numerical study of inelastic scatterings by time-dependent random potentials in two-dimensional systems
T. Nakanishi and T. Ohtsuki 801

Magnetic localization of classical electrons in 2D disordered lattice
E.M. Baskin and M.V. Entin 805

Chapter 12. Bilayers and multilayers

Superlattice and multilayer structures based on ferromagnetic semiconductor (Ga, Mn)As
A. Shen, H. Ohno, F. Matsukura, H.C. Liu, N. Akiba, Y. Sugawara, T. Kuroiwa and Y. Ohno 809

Macroscopic correlation-driven charge transfer between capacitively isolated electron layers
H.C. Manoharan, Y.W. Suen, T.S. Lay, M.B. Santos, M. Shayegan 814

Many-body interactions, the quantum Hall effect, and insulating phases in bilayer two-dimensional hole-gas systems
A.R. Hamilton, M.Y. Simmons, T.G. Griffiths, A.K. Savchenko, M. Pepper and D.A. Ritchie 819

Properties of the soliton-lattice state in double-layer quantum Hall systems
C.B. Hanna, A.H. MacDonald and S.M. Girvin 824

Coherent states in the bilayer fractional quantum Hall ferromagnet
T. Nakajima and H. Aoki 828

Quantum Hall effect transitions in a bilayer electron system in tilted magnetic fields
T.S. Lay, T. Jungwirth, L. Smrčka and M. Shayegan 832

Interlayer quantum coherence and anomalous stability of $\nu = 1$ bilayer quantum Hall state
A. Sawada, Z.F. Ezawa, H. Ohno, Y. Horikoshi, S. Kishimoto, F. Matsukura, Y. Ohno, M.Yasumoto and A. Urayama 836

Quantum coherence in quantum Hall ferromagnet
Z.F. Ezawa 841

Magnetic breakdown and Landau level spectra of a tunable double-quantum-well Fermi surface
J.A. Simmons, N.E. Harff, S.K. Lyo, G.S. Boebinger, J.F. Klem, L.N. Pfeiffer and K.W. West 845

Onset of subband locking in double-quantum-well structures as the signature of wave function delocalization
K.R. Zolleis, C.H.W. Barnes, A.G. Davies, M.Y. Simmons, D.A. Ritchie and M. Pepper 850

Interaction effects on cyclotron resonance in Semiconductor double quantum well structures
H. Sakakibara and K. Hirakawa 855

The frictional drag between coupled two-dimensional electronic systems in magnetic fields
H. Rubel, A. Fischer, W. Dietsche, C. Jörger, K. von Klitzing and K. Eberl 859

Frictional drag mediated by acoustic phonons
M.C. Bønsager, K. Flensberg, B.Y.-K. Hu and A.H. MacDonald 864

Electron–electron scattering between closely spaced two-dimensional electron gases
N.P.R. Hill, J.T. Nicholls, E.H. Linfield, M. Pepper, D.A. Ritchie, B.Y.-K. Hu and K. Flensberg 868

Chapter 13. Supperlattices and quantum wells

Change of dimensionality of semiconductor superlattices induced by an in-plane magnetic field
O. Jaschinski, G. Nachtwei, J. Schoenes, P. Bönsch and A. Schlachetzki 873

Carrier density dependence of transitions between chaotic and periodic photocurrent oscillations in undoped GaAs/AlAs superlattices
N. Ohtani, N. Egami, H.T. Grahn and K.H. Ploog 878

Angular dependent magnetoresistance oscillation in GaAs/AlGaAs superlattice
M. Kawamura, A. Endo, M. Hirasawa, S. Katsumoto and Y. Iye 882

Resonance effect of coherent folded acoustic phonons generated by ultrashort light pulses in GaAs/AlAs superlattice
K. Mizoguchi, K. Matsutani, M. Hase, S. Nakashima and M. Nakayama 887

High-field miniband transport in semiconductor superlattices under the influence of a transverse magnetic field
P. Kleinert 891

Magnetic-field induced metal–insulator transition due to spin effects in Si/SiGe superlattices
A. Prinz, G. Brunthaler, T. Dietl, M. Sawicki, J. Jaroszyński, F. Schäffler and G. Bauer 895

Piezoelectric potential under two-dimensional superlattices
D.E. Petticrew, J.H. Davies, A.R. Long and I.A. Larkin 900

Photon assisted electric field domains in doped semiconductor superlattices
R. Aguado and G. Platero 904

Magneto-photoluminescence spectra of GaP/AlP short-period superlattices in high magnetic fields and uniaxial pressures
K. Uchida, N. Miura, T. Sugita, F. Issiki, N. Usami and Y. Shiraki 909

Novel magneto-resistance oscillations in short period lateral superlattices
C. Albrecht, J.H. Smet, D. Weiss, K. von Klitzing, V. Umansky and H. Schweizer 914

Transitions between fractional and integer quantum Hall states in a lateral superlattice potential
D. Pfannkuche and A.H. MacDonald 918

Quantum effects in the electronic conductivity of lateral superlattices
F. Petit, M. Hayne, F. Lelarge and B. Etienne 922

High-field magneto-spectroscopy of quasi-2D electrons and holes in IAs/$Al_xGa_{1-x}Sb$ single quantum wells
S.K. Singh, B.D. McCombe, Y.J. Wang, J. Kono, I. Lo, W.C. Mitchel and C.E. Stutz 927

Cyclotron resonance in ZnCdSe/ZnSe multi-quantum wells at ultra-high magnetic fields up to 150 T
Y. Imanaka and N. Miura 932

Effect of finite temperature correlations on plasmon and Coulomb drag in coupled quantum wells
L. Liu, L. Świerkowski and D. Neilson 937

Wave-function spectroscopy in parabolic quantum wells
G. Salis, E. Ensslin, K.B. Campman, K. Maranowski and A.C. Gossard 941

Anomalous carrier lifetime enhancement and effective mass discontinuity observed during magnetic-field-induced subband depopulation in a wide parabolic quantum well
G.R. Facer, B.E. Kane, R.G. Clark, L.N. Pfeiffer and K.W. West 946

Magneto-photoluminescence anomalies at integer and fractional quantum Hall regimes in CdTe–CdMgTe modulation n-doped single quantum wells
H. Kunimatsu, S. Takeyama, K. Uchida, N. Miura, G. Karczewski, T. Wojtowicz and J. Kossut 951

Magnetic-field-dependent multi-line cyclotron resonance in modulation-doped GaAs/AlGaAs quantum wells
S.R. Ryu, Z.X. Jiang, T.M. Yeo, B.D. McCombe, Y.J. Wang and W. Schaff 955

Nonlinear and oscillatory electronic transport in superlattices as a probe of structural imperfections
E. Schöll, G. Schwarz and A. Wacker 961

Cyclotron resonance studies of strongly coupled double quantum wells in tilted magnetic fields near the quantum and semi-classical limits
T.P. Marlow, D.D. Arnone, C.L. Foden, E.H. Linfield, D.A. Ritchie and M. Pepper 966

Far-infrared emission and possibility of population inversion of hot holes in MQW InGaAs/GaAs heterostructures under real space transfer
V.Ya. Aleshkin, A.A. Andronov, A.V. Antonov, N.A. Bekin, V.I. Gavrilenko, I.G. Malkina, D.G. Revin, E.A. Uskova and B.N. Zvonkov 971

List of Contributors 976

Subject Index 981